

ARITHMETIC SEQUENCES OF HIGHER DEGREES CHARACTERIZING FIGURATE NUMBERS

Grażyna Rygał^a, Arkadiusz Bryll^b, Grzegorz Bryll^c

^a*Pedagogical Faculty
Jan Długosz University in Częstochowa
Armii Krajowej 13/15, 42-200 Częstochowa, Poland
e-mail: g.rygal@op.pl*

^b*Technical University of Częstochowa, The Faculty of Management
Dąbrowskiego 69, 42-200 Częstochowa, Poland*

^c*Emeritus professor, Poland*

Abstract. Figurate numbers have simple geometric illustration: polygonal numbers can be represented by polygons, pyramidal numbers by pyramids, prismatic numbers by prisms, and trapezoidal numbers by trapezoids. The numbers mentioned above can be defined by formulae¹ or can be characterized by some arithmetic sequences of higher degrees which allow to obtain the corresponding formulae [3]. Figurate numbers due to their geometrical illustration and interesting properties can be of interest for school pupils.

1. Arithmetic sequences of higher degrees

For arbitrary sequence $\{a_n\} : a_1, a_2, a_3, \dots$ we can calculate the sequence of the first finite differences $\{\Delta^1 a_n\}$

$$\Delta^1 a_1 = a_2 - a_1, \quad \Delta^1 a_2 = a_3 - a_2, \quad \Delta^1 a_3 = a_4 - a_3, \dots \quad (1)$$

and the sequences of successive differences

$$\Delta^{k+1} a_i = \Delta^k a_{i+1} - \Delta^k a_i, \quad i = 1, 2, 3, \dots \quad k = 1, 2, 3, \dots \quad (2)$$

Using the method of complete mathematical induction it can be proved that an arbitrary term of a sequence $\{a_n\}$ can be described by the following formula (see [1, 4]):

¹This method was used by W. Sierpiński in the book [5] in definitions of triangle and tetrahedral numbers.

$$a_n = \binom{n-1}{0} a_1 + \binom{n-1}{1} \Delta^1 a_1 + \binom{n-1}{2} \Delta^2 a_1 + \dots + \binom{n-1}{n-1} \Delta^{n-1} a_1. \quad (3)$$

It is evident that to define the n th term of a sequence $\{a_n\}$ it is sufficient to (know) have the first term and the differences: $\Delta^1 a_1, \Delta^2 a_1, \dots, \Delta^{n-1} a_1$.

A sequence $\{a_n\}$ is called *an arithmetic sequence of the degree m* ($m = 1, 2, \dots$) if and only if the sequence $\{\Delta^m a_n\}$ is constant and differs from the zero sequence. A constant sequence will be called an arithmetic sequence of the zero degree.

It follows from Eq. (3) that an arbitrary term of an arithmetic sequence of the degree m is expressed as:

$$a_n = \binom{n-1}{0} a_1 + \binom{n-1}{1} \Delta^1 a_1 + \binom{n-1}{2} \Delta^2 a_1 + \dots + \binom{n-1}{m} \Delta^m a_1. \quad (4)$$

To determine the differences $\Delta^1 a_1, \Delta^2 a_1, \dots, \Delta^m a_1$ we draw up the following table

$$\begin{array}{cccc}
 a_1 & & & \\
 \Delta^1 a_1 & & & \\
 a_2 & \Delta^2 a_1 & & \\
 \Delta^1 a_2 & & \Delta^3 a_1 & \dots \\
 a_3 & \Delta^2 a_2 & & \\
 \Delta^1 a_3 & & \Delta^3 a_2 & \\
 a_4 & \Delta^2 a_3 & & \\
 \Delta^1 a_4 & \vdots & \vdots & \\
 a_5 & \vdots & \vdots & \vdots \\
 \vdots & \vdots & \vdots & \vdots
 \end{array} \quad (5)$$

2. Polygonal numbers

For sequences with general terms

- (a) $a_n = n$,
- (b) $a_n = 2n - 1$,
- (c) $a_n = 3n - 2$,
- (d) $a_n = 4n - 3$,

(e) $a_n = 5n - 4,$

(f) $a_n = (s - 2)n - (s - 3), \quad s = 3, 4, 5, \dots$


the sequences of partial sums² have the following form, respectively:

$\{t^{(3)}(n)\} :$	1	3	6	10	15	21	28	...
$\{t^{(4)}(n)\} :$	1	4	9	16	25	36	49	...
$\{t^{(5)}(n)\} :$	1	5	12	22	35	51	70	...
$\{t^{(6)}(n)\} :$	1	6	15	28	45	66	91	...
$\{t^{(7)}(n)\} :$	1	7	18	34	55	81	112	...
.....
$\{t^{(s)}(n)\} :$	1	s	$3s - 3$	$6s - 8$	$10s - 15$	$15s - 24$	$21s - 35$...


Elements (terms) of a sequence $\{t^{(s)}(n)\}$ are called s -gonal numbers. Hence, there are trigonal numbers, square (quaternal) numbers, pentagonal numbers, hexagonal numbers, etc.

Geometrical illustration of these numbers is as follows:

Trigonal numbers:




Square numbers:




²The n th partial sum for a sequence $\{a_n\}$ is $a_1 + a_2 + \dots + a_n$

Pentagonal numbers:



Hexagonal numbers:



The sequences $\{t^{(s)}(n)\}$ ($s = 3, 4, \dots$) are arithmetic sequences of the second degree. The general term of a sequence $\{t^{(s)}(n)\}$ can be determined by the method of successive differences using table (5) and equation (4). The table of differences for this sequence has the form:

1			
s	$s - 1$	$s - 2$	
$3s - 3$	$2s - 3$	$s - 2$	0
$6s - 8$	$3s - 5$	$s - 2$	0
$10s - 15$	$4s - 7$	\vdots	\vdots
\vdots	\vdots	\vdots	\vdots

As $t^{(s)}(1) = 1$, $\Delta^1 t^{(s)}(1) = s - 1$, $\Delta^2 t^{(s)}(1) = s - 2$, then, using Eq. (4), we obtain:

$$t^{(s)}(n) = \binom{n-1}{0} \cdot 1 + \binom{n-1}{1}(s-1) + \binom{n-1}{2}(s-2),$$

i.e.

$$t^{(s)}(n) = \frac{n}{2}[n(s-2) - s + 4]. \tag{6}$$

Therefore trigonal, square, pentagonal, hexagonal, and heptagonal numbers can be defined using the following equations:

$$\begin{aligned} t^{(3)}(n) &= \frac{n}{2}(n+1), & t^{(4)}(n) &= n^2, \\ t^{(5)}(n) &= \frac{n}{2}(3n-1), & t^{(6)}(n) &= \frac{n}{2}(4n-2), \\ t^{(7)}(n) &= \frac{n}{2}(5n-3). \end{aligned}$$

Many properties of s -gonal numbers can be found in books [2, 5].

3. Pyramidal numbers

If for a sequence $\{t^{(s)}(n)\}$ the sequence of partial sums is created, then a sequence $\{T^{(s)}(n)\}$ is obtained being an arithmetic sequence of the third degree. Elements (terms) of this sequence are called *s-gonal pyramidal numbers*.

The sequences of trigonal, square, pentagonal, hexagonal, heptagonal, ..., s -gonal pyramidal numbers have the following form

$\{T^{(3)}(n)\} :$	1	4	10	20	35	...
$\{T^{(4)}(n)\} :$	1	5	14	30	55	...
$\{T^{(5)}(n)\} :$	1	6	18	40	75	...
$\{T^{(6)}(n)\} :$	1	7	22	50	95	...
$\{T^{(7)}(n)\} :$	1	8	26	60	105	...
.....
$\{T^{(s)}(n)\} :$	1	$s+1$	$4s-2$	$10s-10$	$20s-25$...

respectively.

To determine the general term of a sequence $\{T^{(s)}(n)\}$ draw up a table of successive differences:

$$T^{(5)}(n) = \frac{n^2}{2}(n+1), \quad T^{(6)}(n) = \frac{n}{6}(n+1)(4n-1),$$

$$T^{(7)}(n) = \frac{n}{6} \cdot (n+1)(5n-2).$$

In the book [5] many properties of trigonal pyramidal numbers being elements (terms) of the sequence $T^{(3)}(n)$ can be found.

4. Prismatic numbers

Let m be an arbitrary natural number distinct from 1. Consider the sequence:

$$m, \quad 2(m+1), \quad 3(m+2), \quad 4(m+3), \quad 5(m+4), \dots$$

and create for it the sequence $\{P^{(m)}(n)\}$ being the sequence of partial sums:

$$\{P^{(m)}(n)\} : \quad m, \quad 3m+2, \quad 6m+8, \quad 10m+20, \quad 15m+40, \dots$$

It is easy to see that the above sequence is an arithmetic sequence of the third degree. The general term of this sequence can be found drawing up a table of successive differences

	m			
		$2m+2$		
$3m+2$			$m+4$	
		$3m+6$		2
$6m+8$			$m+6$	0
		$4m+12$		2
$10m+20$			$m+8$	0
		$5m+20$		2
				\vdots
$15m+40$			$m+10$	\vdots
		$6m+30$		\vdots
				\vdots
$21m+70$				\vdots
				\vdots
				\vdots

Then we have

$$P^{(m)}(1) = m, \quad \Delta^1 P^{(m)}(1) = 2m+2,$$

$$\Delta^2 P^{(m)}(1) = m+4, \quad \Delta^3 P^{(m)}(1) = 2.$$

Using Eq. (4), we obtain

$$P^{(m)}(n) = \binom{n-1}{0}m + \binom{n-1}{1}2(m+1) + \binom{n-1}{2}(m+4) + \binom{n-1}{3} \cdot 2$$

or

$$P^{(m)}(n) = \frac{n}{6}(n+1)[3m+2(n-1)]. \quad (9)$$

Elements (terms) of the sequence $\{P^{(m)}(n)\}$ are called *prismatic numbers of the range m* ($m \geq 2$.)

The sequences of prismatic numbers of the second, third, and fourth range have the form:


$$\{P^{(2)}(n)\} : 2 \quad 8 \quad 20 \quad 40 \quad 70 \quad \dots \quad P^{(2)}(n) = \frac{1}{3}(n+1)(n+2)$$

$$\{P^{(3)}(n)\} : 3 \quad 11 \quad 26 \quad 50 \quad 85 \quad \dots \quad P^{(3)}(n) = \frac{n}{6}(n+1)(2n+7)$$


$$\{P^{(4)}(n)\} : 4 \quad 16 \quad 32 \quad 60 \quad 100 \quad \dots \quad P^{(4)}(n) = \frac{n}{3}(n+1)(n+5)$$

Using prism, the prismatic numbers have the following geometrical illustration:

$$m = 2$$



$m = 3$



It should be noted that faces of a prism illustrating prismatic numbers being triangles can be interpreted as corresponding triangular numbers, whereas lateral faces of a prism being trapezes as *trapezoid numbers*.

For numbers $m = 2, m = 3, m = 4, \dots, m = s$ we obtain the following sequences of trapezoid numbers:

$$\begin{array}{l}
 \{R^{(2)}(n)\} : \quad 2 \quad 5 \quad 9 \quad 14 \quad 20 \quad \dots \\
 \{R^{(3)}(n)\} : \quad 3 \quad 7 \quad 12 \quad 18 \quad 25 \quad \dots \\
 \{R^{(4)}(n)\} : \quad 4 \quad 9 \quad 15 \quad 22 \quad 30 \quad \dots \\
 \dots \dots \dots \dots \dots \dots \dots \dots \\
 \{R^{(s)}(n)\} : \quad s \quad 2s + 1 \quad 3s + 3 \quad 4s + 6 \quad 5s + 10 \quad \dots
 \end{array}$$

The general term of the sequence $\{R^{(s)}(n)\}$ can be found based on a table of successive differences:

s	$s + 1$	
$2s + 1$	$s + 2$	1
$3s + 3$	$s + 3$	0
$4s + 6$	$s + 4$	1
$5s + 10$	\vdots	\vdots
\vdots	\vdots	\vdots

Hence, we have $R^{(s)}(1) = s$, $\Delta^1 R^{(s)}(1) = s + 1$, $\Delta^2 R^{(s)}(1) = 1$. Using Eq. (4), we obtain

$$R^{(s)}(n) = \binom{n-1}{0} s + \binom{n-1}{1} (s+1) + \binom{n-1}{2}$$

or

$$R^{(s)}(n) = \frac{n}{2}(n-1+2s). \quad (10)$$

The sequence $\{R^{(s)}(n)\}$ is an arithmetic sequence of the second degree.

According to (10), the sequences $\{R^{(2)}(n)\}$, $\{R^{(3)}(n)\}$, $\{R^{(4)}(n)\}$ have the following general terms:

$$R^{(2)}(n) = \frac{n}{2}(n+3), \quad R^{(3)}(n) = \frac{n}{2}(n+5), \quad R^{(4)}(n) = \frac{n}{2}(n+7).$$

Summarizing we state that all the types of the abovementioned figurate numbers can be characterized by some arithmetic sequences of the second or third degree.

References

- [1] G. Bryll, R. Sochacki. *Wybrane zagadnienia dydaktyki matematyki*. Oficyna Wydawnicza GARMOND, Poznań 2009.
- [2] Sz. Jeleński. *Śladami Pitagorasa*. PZWS, Warszawa 1953.
- [3] H.B. Lübsen. *Analysis*. Friedrich Brandstetter Verlag, Leipzig 1922.
- [4] A. Mostowski, M. Stark. *Elementy algebry wyższej*. PWN, Warszawa 1958.
- [5] W. Sierpiński. *Liczby trójkątne*. Biblioteczka Matematyczna Nr 12, PZWS, Warszawa 1962.